

**Working paper 8/2013****A contest success function for ranking****Alberto Vesperoni, University of Siegen**

Abstract: *A contest is a game where several players compete for winning prizes by expending costly efforts. We assume that the outcome of a contest is an ordered partition of the set of players (a ranking) and a contest success function assigns a probability to each possible outcome as a function of players' efforts. We define a contest success function for contests whose outcome is a ranking of any type, i.e., with any number of players at each rank. This approach is new in contest theory since the axiomatic work has exclusively been on contests with single-winner, whose outcome is a ranking with one player in the first rank and all other players in the second rank. The contest success function is characterized by pair-swap consistency, which is an axiom of independence of irrelevant alternatives and generalizes the main axiom in Skaperdas (1996).*

A contest success function for rankings

Alberto Vesperoni*

November 2013

Abstract

A contest is a game where several players compete for winning prizes by expending costly efforts. We assume that the outcome of a contest is an ordered partition of the set of players (a *ranking*) and a contest success function assigns a probability to each possible outcome as a function of players' efforts. We define a contest success function for contests whose outcome is a ranking of any type, i.e., with any number of players at each rank. This approach is new in contest theory since the axiomatic work has exclusively been on contests with single-winner, whose outcome is a ranking with one player in the first rank and all other players in the second rank. The contest success function is characterized by *pair-swap consistency*, which is an axiom of independence of irrelevant alternatives and generalizes the main axiom in Skaperdas (1996).

Keywords: conflict, contest, ranking, success function, axiom, probabilistic choice.

JEL classification: C70, D72, D74, C25, D81.

1 Introduction

A contest is a game where several players compete for winning prizes by expending costly efforts. Contest models are used to study situations of conflict such as war, political lobbying, litigation and R&D races. In this paper we generalize the framework and develop a model for studying a broader class of competitive environments.

In the literature so far, the axiomatic work exclusively focused on contests with single winner, where one player wins the prize while all others lose. However, it seems particularly natural that a contest may induce different types of outcome. For instance, in the academic job market candidates compete for various research and teaching positions and most of them are recruited for some job. In the competition for tuition waiver rights at a university, some percentage of best performing students may be awarded a full tuition waiver, some others half and the remaining nothing. In both these examples, the outcome of the contest is different than the single-winner type. In this paper we define a model for contests with any type of outcome and characterize a class of contests by a property of independence which, loosely speaking, requires the relative performance of any two players in a contest to depend only on their own relative efforts, and not on the efforts of others.

*FoKoS Institute, University of Siegen, alberto.vesperoni@gmail.com

Let us describe the model in more detail. We assume that the outcome of a contest is a *ranking*, which is defined as an ordered partition of the set of players, assigning a rank (or *level*) to each player. The type of a ranking is identified by the exact number of players at each level. When a ranking is *strict*, each player has a different level and no level is shared by more than one player. When a ranking is *non-strict*, at least one level is shared by more than one player. There are many types of non-strict ranking. The outcome of a single-winner contest is a non-strict ranking where one player is ranked first and all other players are ranked second. The awarding of tuition waiver rights at a university induces a non-strict ranking of students, dividing them into three categories. The analysis here allows for the study of any type of ranking as the outcome of a contest.

The outcome of a contest is stochastically determined by players' efforts. More precisely, each profile of efforts induces a non-degenerate probability distribution over all possible outcomes of a contest. In the example of the awarding of tuition waiver rights, the more each student has prepared during the academic year the higher its probability of being awarded a tuition waiver. This probability is always positive, as even a completely unprepared student has some chances of passing exams. The function which maps players' efforts into probabilities of outcomes is called *contest success function*.

In the literature so far, the axiomatic characterizations of contest success functions are exclusively for contests with single winner. In this paper we axiomatically characterize a contest success function for contests with any type of outcome. We characterize our success function by a unique property, *pair-swap consistency*. Pair-swap consistency is an axiom of independence, requiring the relative performance of two players in a contest of any type to be exclusively determined by their own efforts, as if they were the only players in the contest. More precisely this axiom requires, for each pair of players, the probability of any ranking where one of them is one level above the other to be proportional to the probability of the same player winning against the other in a two-player contest.

Pair-swap consistent contest success functions are appropriate for modeling certain situations more than others. They are appropriate for modeling competitive situations where each player acts individually and fights each opponent indiscriminately. The academic job market and the awarding of tuition waiver rights at a university are reasonable examples of such cases. On the other hand, these functions are inappropriate for modeling situations where players compete unevenly against their opponents. Examples of such situations are when some players collude, or when there are geographical or cultural barriers to competition between some but not all players.

1.1 Related models in contest theory and probabilistic choice

Restricting our attention to contests of single-winner type, our contest success function is equivalent to the one characterized in Skaperdas (1996), the most well known in the literature. Our axiom is analogous to *sub-contest consistency*, which is the main axiom in Skaperdas (1996) and in most axiomatic characterizations of contest success functions. Loosely speaking, sub-contest consistency is a property of success functions for contests of single-winner type which requires the winning probability of a player to be proportional to its winning probability in sub-contests restricted to *any subset* of players. For contests of this type, pair-swap consistency is equivalent to a weaker version of sub-contest consistency, requiring the property to hold only for sub-contests restricted to *any pair* of players.

Our framework is also meaningful in probabilistic choice and random utility theory. A contest among n players can be interpreted as the probabilistic preferences of a decision maker over a set of n alternatives. Efforts of players correspond to underlying qualities of alternatives and the contest success function gives the probability distribution of the preferences. Pair-swap consistency is analogous to Luce (1959)'s *choice axiom*, also known as *independence of irrelevant alternatives*. This axiom considers probabilistic choices from different subsets of alternatives and, roughly speaking, it requires the relative probability of any two alternatives to be independent of other alternatives in the subset. Independence of irrelevant alternatives characterizes an important class of probabilistic choice functions, which includes the *multinomial logit*. The framework developed here generalizes Luce (1959), in that it allows to represent not only probabilistic choices but probabilistic preferences of any type.¹

1.2 Applications in contest theory

In the last section of this paper we apply our success function to contest games with different structures. More specifically, we analyze equilibrium behavior in contests with various types of outcome, number of stages, number and values of prizes, in order to identify which of these structures induces the highest equilibrium efforts. These topics have been widely studied with success functions of perfectly discriminating type.² However, the study of these subjects with success functions of imperfectly discriminating type is still incomplete.³ Our approach provides a framework for generalizing the analysis, as it allows to model any of these structures.

The paper develops as follows. Section 2 reviews the related literature in contest theory and probabilistic choice. Section 3 describes the model, and Section 4 defines the success function and pair-swap consistency, characterizing the success function via this axiom. Section 5 analyzes some properties of the success function and studies equilibrium behavior in contest games with various structures.

2 Related literature

The axiomatic approach in contest theory started with the seminal work of Skaperdas (1996), which defines a contest success function for single-winner contests. The function is characterized by five axioms: probability, monotonicity, anonymity, sub-contest independence and sub-contest consistency.⁴ While the first three axioms are straightforward, the last two are

¹Preferences may be of various non-strict types as well as strict due to the decision maker's bounded rationality. More specifically, a decision maker may be unable to process all necessary information for ordering the alternatives in a strict ranking.

²A success function is of perfectly discriminating type if the outcome of a contest is non-stochastically determined by players' efforts. For example in an auction players are deterministically ranked according to their bids. Baye et al. (1993, 1996), Clark and Riis (1998a), Moldovanu and Sela (2001, 2006) and Konrad and Kovenock (2009) study equilibrium behavior in perfectly discriminating contests with various structures.

³A contest success function is of imperfectly discriminating type if efforts of players induce a non-degenerate probability distribution over all possible outcomes of a contest. Clark and Riis (1996, 1998c), Wärneryd (2001) and Fu and Lu (2009, 2012) analyze imperfectly discriminating contests with different structures.

⁴These terms are not used in Skaperdas (1996). We follow the terminologies in Wärneryd (2001) and Münster (2009).

not. Let us briefly describe them. Given a contest among some players, a sub-contest is a contest restricted to a subset of these players. *Sub-contest independence* requires the players' winning probabilities in any sub-contest to be independent of the efforts of players which are excluded from the sub-contest. *Sub-contest consistency* demands a player's winning probability to be proportional to its winning probability in any sub-contest. More specifically, given a set of players N , for any player $i \in N$ and any subset of players $M \subset N$ where $i \in M$, sub-contest consistency requires that player i 's winning probability $p(i)$ satisfies $p(i)/\left[\sum_{j \in M} p(j)\right] = p_M(i)$, where $p_M(i)$ is player i 's winning probability in the sub-contest restricted to players in M . While the other axioms are fairly weak, sub-contest consistency is clearly demanding. In this sense we say that sub-contest consistency is the *main* axiom of the characterization. In Skaperdas (1996), a contest success function p fulfills these five axioms if and only if there exists an increasing function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that each player $i \in N$'s winning probability can be represented as $p(i) = f(x_i)/\left[\sum_{j \in N} f(x_j)\right]$ if $\sum_{j \in N} f(x_j) > 0$ and $1/n$ otherwise, where $x_j \geq 0$ is the effort of player $j \in N$ and $n \geq 2$ is the number of players in N .

The contest success function defined in Skaperdas (1996) has been generalized in several directions by characterizations of different success functions. In all these extensions the outcome of a contest is of single-winner type, and most characterizations relax some of the first four axioms in Skaperdas (1996) and maintain sub-contest consistency. Let us briefly discuss them. Blavatsky (2010) relaxes probability, introducing a success function for modeling contests with the possibility of a draw. Clark and Riis (1998b) relax anonymity, developing a success function which treats players unfairly. Rai and Sarin (2009) and Arbatskaya and Mialon (2010) define success functions for multi-dimensional efforts of players, and Münster (2009) axiomatizes a success function for contests between groups of players. These last three extensions generalize the framework in Skaperdas (1996) to cases where a player's effort is a vector instead of a scalar⁵, and their axioms are generalizations of the ones in Skaperdas (1996) to the multi-dimensional case. Bozbay and Vesperoni (2013) develop a success function for contests where each pair of players can be ally or enemy of each other, defining the function for any possible network of alliances. In their characterization, most axioms are violated for any network where some players are allies.

This paper is first in axiomatically characterizing a contest success function for rankings different than the single-winner type. However, there is non-axiomatic work on the subject. Clark and Riis (1996) introduce a success function for contests whose outcome is a strict ranking of players, without providing an axiomatic characterization.⁶ Fu and Lu (2011) show that this contest success function is the outcome of a stochastic process, under certain assumptions on the distribution of errors.⁷ Fu et al. (2013) propose an alternative success function for contests whose outcome is a strict ranking, deriving it from a different stochastic

⁵In Münster (2009), a group of $n \geq 2$ players can be interpreted as a single player with n dimensional effort.

⁶Clark and Riis (1996) derive their success function for strict-ranking contests from a sequential procedure. More precisely, they define a strict-ranking contest as a grand contest with multiple stages, where in each stage a single-winner contest with success function of Skaperdas (1996) takes place. All players participate in the first stage's contest, whose unique winner is the player ranked first in the grand contest. All players participate in the second stage except the first stage's winner. The winner of the second stage's contest is the player ranked second in the grand contest. In a similar way, the sequential procedure orders all players in a strict ranking.

⁷The crucial assumption of their characterization is the independence of the errors of each stage of the sequential procedure in Clark and Riis (1996).

process.⁸ The success function developed here is more general than Clark and Riis (1996) and Fu et al. (2013) in that, besides the possibility of being applied to contests whose outcome is a strict ranking, it proposes a unified framework for modeling contests whose outcome is of any type, including non-strict rankings.

The second body of literature that this paper connects to is probabilistic choice and random utility theory. Luce (1959) axiomatically characterizes a framework for modeling the probabilistic choice of a decision maker from a set of alternatives. The main axiom of the characterization is *independence of irrelevant alternatives* (IIA). McFadden and Zarembka (1974) axiomatize the most popular functional form of this model, known as the *multinomial logit*. McFadden (1973) provides a non-axiomatic characterization, showing that the choice probabilities can be derived from a stochastic process with errors of the *generalized extreme value* (GEV) family, where the crucial assumption is the independence of these errors. By allowing for correlation of errors, Williams (1977) and McFadden (1978) generalize the framework into a model known as *nested logit*. Beggs et al. (1981) generalize the multinomial logit model in a different direction, providing a representation of probabilities of *strict rankings* of alternatives known as *rank-ordered logit*. Their characterization is non-axiomatic and generalizes the one in McFadden (1973), showing that their model can be derived from a stochastic process with errors of GEV type. While their representation is only for *strict* rankings, here we provide a model for any type of rankings, and our characterization generalizes Luce (1959).

3 Model

There is a set of players $N = \{1, \dots, n\}$, where $n \geq 2$. A *ranking* r is an ordered partition of N which assigns a level to each player. We write $r(i) = l$ to say that in ranking r player i is ranked at level l . For $N = \{1, 2, 3\}$, if $r(1) = 1$, $r(2) = 1$ and $r(3) = 2$, we write $r = (1, 2; 3)$. We exclude the degenerate case where all players are ranked equally.⁹ Then, the set of all rankings R is the set of all ordered partitions of N except (N) . We say that player i is *ranked above* player j if $r(i) < r(j)$.

A *type of ranking* describes the exact number of players in each level of a ranking. We write $t(l) = m$ to say that a ranking of type t assigns m players to level l .¹⁰ For instance, any ranking of single-winner type has one player at the first level ($t(1) = 1$) and all other players at the second level ($t(2) = n - 1$). We denote by T the set of all types of ranking.

Given a type of ranking $t \in T$, the *set of possible outcomes* of a contest is the set of all rankings in R of the same type t .¹¹ For instance, the set of possible outcomes of a single-

⁸Fu et al. (2013) derive this success function by reversing the sequential procedure in Clark and Riis (1996). In other words, instead of starting by assigning the best performing player to the first level of the ranking, they start by assigning the worst performing player to the last level.

⁹Formally, a function $r : N \rightarrow N$ represents an ordered partition of N if and only if no player is assigned to a level below some empty level (for any i and l in N we have $r(i) = l$ only if for each $l' < l$ there exists $j \in N$ such that $r(j) = l'$). A function r which fulfills this condition is a ranking if and only if there is no level to which all players belong ($r(i) \neq r(j)$ for some $i, j \in N$).

¹⁰More formally, a function $t : N \rightarrow N$ is a type of ranking if and only if (1) each player is assigned to some level ($\sum_{l \in N} t(l) = n$); (2) there is no level to which all players belong ($t(l) \leq n - 1$ for all $l \in N$); (3) if a level has some players in it then there are no empty levels above that level ($t(l) \geq 1$ only if $t(l') \geq 1$ for all $l' < l$ and $l \geq 2$).

¹¹A set of rankings $A \subseteq R$ is a set of possible outcomes if and only if (1) every $r \in A$ is of same type $t \in T$; (2) for each $r \in A$, there exists $r' \in A$ such that $r'(\psi(i)) = r(i)$ for any player $i \in N$ and permutation

winner contest is the set of all rankings in R of single-winner type. We denote by $R(t)$ the set of possible outcomes of type $t \in T$, and by \mathcal{R} the set of all sets of possible outcomes.

Each player $i \in N$ is associated with an effort $x_i \in X_i \subseteq \mathbb{R}_+$, and $x \in X \subseteq \mathbb{R}_+^n$ is an effort profile. Given any type of ranking $t \in T$, a *contest success function* defines the probability of each possible outcome $r \in R(t)$ for any effort profile $x \in X$. Then, for any type $t \in T$, a contest success function is a mapping $p : R(t) \times X \rightarrow [0, 1]$ where $\sum_{r \in R(t)} p(r, x) = 1$ for every $x \in X$, and we write $p(r, x)$ for the probability of $r \in R(t)$ being the outcome for every $x \in X$.

4 Characterization of the contest success function

Let us define a contest success function for any type of ranking and characterize it via *pair-swap consistency*. To define this axiom, we first define a *pair-swap ranking*. Let $t \in T$ be any type. For each ranking $r \in R(t)$ and each pair of players $i, j \in N$, the pair-swap ranking $r_{i,j}$ is obtained from the ranking r by players i and j swapping their levels and all other players remaining at the same levels.¹² For instance, for $N = \{1, 2, 3, 4\}$, if $r = (2; 3; 4; 1)$ the ranking $r_{2,4}$ is defined as $(4; 3; 2; 1)$, where players 2 and 4 swap their levels and players 3 and 1 remain at the same levels as in r . We can now define pair-swap consistency.

Definition 4.1 *Given a function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_{++}$ and a type of ranking $t \in T$, a contest success function satisfies **pair-swap consistency** if for each effort profile $x \in X$, ranking $r \in R(t)$ and pair of players $i, j \in N$ with $r(j) = r(i) + 1$*

$$\frac{p(r, x)}{p(r, x) + p(r_{i,j}, x)} = \frac{f(x_i)}{f(x_i) + f(x_j)}.$$

For each pair of players $i, j \in N$ and effort profile $x \in X$, the RHS of the equation in Definition 4.1 is the winning probability of player i in a two-player contest against player j , where the success function of the two-player contest belongs to the family characterized in Skaperdas (1996).¹³ For the same effort profile $x \in X$ and any type of ranking $t \in T$, the LHS is the probability of a ranking $r \in R(t)$ where player i is one level above player j , conditional on all players in $N \setminus \{i, j\}$ being at given levels in the outcome.¹⁴ In other words, the LHS measures how likely it is that i is ranked above j rather than j above i *everything else is equal*, i.e., when any other player is at a given level. Then, we can say that pair-swap consistency defines a property of a contest success function $p : R(t) \times X \rightarrow [0, 1]$ for any type of ranking $t \in T$ which requires, everything else equal, the relative performance of any

ψ of N .

¹²Given a type $t \in T$, for any ranking $r \in R(t)$ and pair of players $i, j \in N$ the pair-swap ranking $r_{i,j}$ is defined as the ranking in $R(t)$ where (1) $r_{i,j}(j) = r(i)$ and $r_{i,j}(i) = r(j)$; (2) $\forall k \in N \setminus \{i, j\} : r_{i,j}(k) = r(k)$.

¹³In Skaperdas (1996) the function f is weakly positive, while here f is required to be *strictly* positive. This is to avoid well known problems in the characterization in Skaperdas (1996) which arise when $f(x_i) = 0$ for some $i \in N$ and $x_i \in X_i$. In short, the contest success function in Skaperdas (1996) is not well defined when this is the case. Moreover in Skaperdas (1996) the function f is *increasing*, while we do not impose such restriction here. We do this only to keep the model more general, as one can imagine competitive situations where a player's performance is not necessarily monotonic in its own effort.

¹⁴Given an effort profile $x \in X$, the outcome of a contest is a random variable. Let us denote it by ω , where $\omega(i)$ is the level of a player $i \in N$ in the outcome. Then, ω takes value in $R(t)$ and its probability distribution is determined by $\Pr(\omega = r) = p(r, x)$ for any $r \in R(t)$. The LHS of the equation in Definition 4.1 can be written as $\Pr(\omega = r | \omega(h) = r(h) \text{ for any } h \notin \{i, j\})$.

pair of players to depend only on their relative efforts, as if they were the only players in the contest. Theorem 4.1 defines the contest success function.

Theorem 4.1 *Given a function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_{++}$ and a type of ranking $t \in T$, a contest success function is pair-swap consistent if and only if for all $r \in R(t)$ and $x \in X$*

$$p(r, x) = \frac{\prod_{i \in N} f(x_i)^{-r(i)}}{\sum_{r' \in R(t)} \prod_{i \in N} f(x_i)^{-r'(i)}}. \quad (1)$$

Proof: See Appendix.

Restricting our attention to contests of single-winner type, the contest success function in Theorem 4.1 is equivalent to the one in Skaperdas (1996). Moreover, for single-winner contests, pair-swap consistency is equivalent to a *weaker version* of sub-contest consistency, which is the main axiom in Skaperdas (1996). While sub-contest consistency requires a player's winning probability in a contest to be proportional to its probability of winning in every sub-contest restricted to any *subset* of players, pair-swap consistency requires this to hold only for sub-contests restricted to any *pair* of players. There is a methodological difference between our characterization and the one in Skaperdas (1996). While we define our success function for a *given* function f , Skaperdas (1996) characterizes a *family* of success functions parametrized by f .¹⁵ Formally, in Theorem 4.1 we define a *consistent extension* of each success function in the family of Skaperdas (1996) to the domain of success functions for any type of ranking.¹⁶ We do so to simplify exposition, as a direct characterization of our function would require heavier notation and repetition of arguments already given in Skaperdas (1996).

5 Discussion

In this Section we study properties of the contest success function defined in Theorem 4.1. We already know that, given a type $t \in T$, the contest success function is a mapping $p : R(t) \times X \rightarrow [0, 1]$ such that $\sum_{r \in R(t)} p(r, x) = 1$ for every $x \in X$. Moreover, given a function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_{++}$, we know that $p(r, x)$ must fulfill pair-swap consistency for any $r \in R(t)$ and $x \in X$. Can we say anything more about our success function? To do so, we additionally assume f to be increasing, weakly concave and twice differentiable.¹⁷ In the following Sections we study properties which hold as a consequence of these.

In Section 5.1 we analyze aspects of the success function which are relevant for studying existence and uniqueness of equilibrium in contest games, i.e., monotonicity and concavity of

¹⁵While in Definition 4.1 pair-swap consistency is defined for a given function f , the axioms in Skaperdas (1996) describe properties of a success function without assuming any functional form. Loosely speaking, Skaperdas (1996) shows that a success function fulfills some axioms if and only if there exists a function f such that each player's winning probability can be represented as a particular function of $(f(x_i))_{i \in N}$.

¹⁶See Thomson (2011) for an introduction to consistent extensions of allocation rules to larger domains. Let us describe our methodology in more detail. We start by choosing a function f and we take the success function corresponding to f from the family in Skaperdas (1996). We use this success function to define pair-swap consistency, which characterizes our success function for a given f and for all types of ranking. Taking any f , we define a family of success functions which fulfills pair-swap consistency.

¹⁷These assumptions are widely accepted in contest theory. See Skaperdas (1996) for axiomatic characterizations of specific functional forms for f which are positive, increasing and twice differentiable. In contest theory the function f is usually assumed to be weakly concave to guarantee the concavity of players' payoffs.

rankings' probabilities in players' efforts.¹⁸ The results show that, as rankings' probabilities are not necessarily monotonic and concave, characterizing an equilibrium is not always trivial. On the other hand this proposition provides a very intuitive framework to reason on the issue. In Section 5.2 we compare probabilities of a player being ranked first in contests with different sets of possible outcomes and same effort profile. These probabilities are generally different from each other and their relative magnitude depends on the effort profile. Our results give insights on how a contest's type affects its precision in ranking the players with the highest efforts at top levels.¹⁹ In Section 5.3 we analyze players' incentives in spending costly efforts in contest games. More precisely, we analyze equilibrium behavior in contests with various structures such as multiple stages, multiple prizes and various types of outcome. Our results highlight which structures maximize equilibrium efforts.

In these Sections we will often refer to contests of three types: single-winner, multi-winner and strict-ranking. Although their meanings are intuitive, let us define them here. For any set of players N , a *single-winner* contest partitions this set such that one player is ranked first ($t(1) = 1$) and all other players are ranked second ($t(2) = n - 1$). A *multi-winner* contest locates $m \geq 2$ players at the first level ($t(1) = m$) and all others at the second ($t(2) = n - m$). In a *strict-ranking* contest the outcome is a strict ranking of players ($t(l) = 1$ for all $l \in N$). Through all the Section we will often refer to the following examples with three players. Consider a single winner-contest with $n = 3$. The set of possible outcomes is $R(t) = \{(1; 2, 3), (2; 1, 3), (3; 1, 2)\}$. Given an effort profile $x \in X$ the probability of ranking $r = (1; 2, 3)$ can be written as

$$p(r, x) = \frac{f(x_1)}{f(x_1) + f(x_2) + f(x_3)} .$$

In a multiple-winner contest with three players and two winners the set of possible outcomes is $R(t) = \{(1, 2; 3), (1, 3; 2), (2, 3; 1)\}$ and for any $x \in X$ the probability of ranking $r = (1, 2; 3)$ is

$$p(r, x) = \frac{f(x_1)f(x_2)}{f(x_1)f(x_2) + f(x_1)f(x_3) + f(x_2)f(x_3)} .$$

Consider a strict-ranking contest with $n = 3$. The set of possible outcomes is

$$R(t) = \{(1; 2; 3), (1; 3; 2), (2; 1; 3), (2; 3; 1), (3; 1; 2), (3; 2; 1)\} ,$$

and for any $x \in X$ the probability of $r = (1; 2; 3)$ takes the value

$$p(r, x) = \frac{f(x_1)^2 f(x_2)}{f(x_1)^2 f(x_2) + f(x_1)^2 f(x_3) + f(x_2)^2 f(x_1) + f(x_2)^2 f(x_3) + f(x_3)^2 f(x_1) + f(x_3)^2 f(x_2)} .$$

¹⁸In single-winner contests, the concavity of f is a sufficient condition for concavity of players's payoffs and for the existence of an equilibrium in pure strategies. See Okuguchi and Szidarovszky (1997), Cornes and Hartley (2005), Esteban and Ray (1999) and Vesperoni (2013) for studies of existence and uniqueness of equilibrium in imperfectly discriminating contests of single-winner type with concave payoffs. If the function f is convex, players' payoffs are not always concave and an equilibrium is typically in mixed strategies. See Baye et al. (1994) and Che and Gale (2000) for analysis of existence and uniqueness of equilibrium in imperfectly discriminating contests of single-winner type where payoffs are not always concave. When f is convex, players' payoffs are similar to the ones in a perfectly discriminating contest. See Baye et al. (1996), Amann and Leininger (1996), Krishna and Morgan (1997) and Siegel (2009) for analysis of existence and uniqueness of equilibrium in perfectly discriminating contests.

¹⁹Perhaps the most interesting applications of this framework are in probabilistic choice and random utility theory. A decision maker wants to choose the best alternative from a set. Its preferences are non strict, due to its bounded rationality in ordering alternatives. This framework allows to study how preference's types affect a decision maker's chances of choosing the best alternative from a set.

5.1 Monotonicity and concavity of rankings' probabilities

In this Section we study rankings' probabilities as functions of players' efforts, and find conditions for their monotonicity and concavity. Let us define some concepts to be used to state our next result. Consider a contest of any type $t \in T$. For any effort profile $x \in X$ and ranking $r \in R(t)$, we say that $p(r, x)$ is *increasing* in the effort of player $i \in N$ if the first derivative $\partial [p(r, x)]/\partial x_i$ evaluated at x is positive, while we say that $p(r, x)$ is *concave* in player i 's effort if the second derivative $\partial^2 [p(r, x)]/\partial x_i^2$ evaluated at x is negative. Before stating our results, we introduce some more notation. For any effort profile $x \in X$, the level of a player $i \in N$ in the outcome is a random variable and its *expected value*²⁰ is $\rho(i, x) := \sum_{r \in R(t)} p(r, x)r(i)$. For each ranking $r \in R(t)$, the *distance* of player i 's level in r from its expected value is $d_i(r, x) := (r(i) - \rho(i, x))^2$, while its *expected distance*²¹ is $\delta_i(x) := \sum_{r \in R(t)} p(r, x)(r(i) - \rho(i, x))^2$.

Proposition 5.1 *For any type $t \in T$ and effort profile $x \in X$, the probability $p(r, x)$ of a ranking $r \in R(t)$ is*

1. *increasing (decreasing) in the effort of a player ranked first (last).*
2. *concave (convex) in the effort of a player ranked first (last) if the type t has only two levels.*
3. *increasing (decreasing) in the effort of a player $i \in N$ if and only if the effort profile x is such that player i 's expected level $\rho(i, x)$ is larger (smaller) than its level $r(i)$.*
4. *concave (convex) in the effort of a player $i \in N$ if the effort profile x is such that player i 's expected level $\rho(i, x)$ is larger (smaller) than its level $r(i)$ and if its expected distance $\delta_i(x)$ is larger (smaller) than its distance $d_i(r, x)$.*

Proof: See Appendix.

Let us briefly interpret the results. Points 1 and 2 of Proposition 5.1 are intuitive, being straightforward generalizations of properties of success functions of single-winner type. The other results are less obvious. Point 3 provides necessary and sufficient conditions for monotonicity of $p(r, x)$. The result follows from the fact that an increase in a player's effort improves its chances of being ranked at better levels. For example, let $n = 4$ and consider any symmetric effort profile $x \in X$. Player $i \in N$'s expected level is $\rho(i, x) = 5/2$. The probability of $r = (1; 2; 3; 4)$ at x is increasing in x_1 and x_2 and decreasing in x_3 and x_4 . Point 4 provides sufficient conditions for concavity of $p(r, x)$. The result can be interpreted as the probability of a ranking presenting decreasing marginal returns if a player's level in the ranking is relatively close to its expected level, and increasing marginal returns otherwise. Consider the example with $n = 4$ and any symmetric $x \in X$. Player $i \in N$'s expected level is $\rho(i, x) = 5/2$ and its expected distance is $\delta_i(x) = 5/4$. The probability of $r = (1; 2; 3; 4)$ at x is concave in x_2 and convex in x_4 .

²⁰The expected level of a player is a function of all players' efforts. Intuitively, a player's expected level increases in its own effort and decreases in others' efforts.

²¹A player's distance and expected distance are functions of all players' efforts. Notice that a player's expected distance is the *variance* of its level in the outcome, hence a measure of dispersion. Then, a player's expected distance is minimized when its expected level is 1 or n , hence when its effort is much larger or much smaller than the others. Conversely, a player's expected distance is maximized when its expected level is the average level, hence at symmetric effort profiles.

Let us see how Proposition 5.1 applies to our examples with three players. Consider the single-winner contest. The probability of ranking $r = (1; 2, 3)$ is increasing and concave in the effort of player 1 and decreasing and convex in the efforts of players 2 and 3 for any $x \in X$. For the multi-winner contest, the probability of $r = (1, 2; 3)$ is increasing and concave in the efforts of players 1 and 2 and decreasing and convex in the effort of player 3 for any $x \in X$. Consider the strict-ranking contest. The probability of $r = (1; 2; 3)$ is increasing in the effort of player 1 and decreasing in the effort of player 3 for any $x \in X$. On the other hand, $p(r, x)$ is increasing (decreasing) in the effort of player 2 if and only if player 2's expected level in the outcome $\rho(2, x)$ is larger (smaller) than its level $r(2) = 2$. Notice that the probability of ranking $r = (1; 2; 3)$ is not necessarily concave in the effort of player 1. One can show that a sufficient condition for its concavity in x_1 is that the expected level is smaller or equal than 2, i.e., $\rho(1, x) \leq 2$.

5.2 Winning probabilities in contests of different types

In this Section we compare a player's probability of being ranked first in a contest of single winner type with its probability of being ranked first in a contest of different type and same effort profile. For any $x \in X$, player $i \in N$'s *probability of being ranked first* in a contest of type $t \in T$ is the joint probability of all outcomes $r \in R(t)$ where this is the case. Then, player i 's probability of being ranked first is $\sum_{r \in R_{i,1}(t)} p(r, x)$, where $R_{i,1}(t)$ is the set of all rankings $r \in R(t)$ where $r(i) = 1$.

Proposition 5.2 *For any number of players $n \geq 3$ and effort profile $x \in X$, a player $i \in N$'s probability of being ranked first in a single-winner contest is*

1. *lower than the probability of being ranked first in a multi-winner contest with same effort profile x .*
2. *higher (lower) than the probability of being ranked first in a contest of a type with one player at first level, one player at second level and $n - 2$ players at third level and same effort profile x if and only if the effort profile x is such that $f(x_i)$ is lower (higher) than*

$$\frac{\sum_{j \neq i} f(x_j)^2 (\sum_{k \neq i, j} f(x_k))}{\sum_{j \neq i} f(x_j) (\sum_{k \neq i, j} f(x_k))}.$$

Proof: See Appendix.

Although the results in Proposition 5.2 are far from being a complete analysis of the subject, they give some idea of the forces at play. The intuition for Point 1 is straightforward. It is easier for any player to be ranked first when there are multiple first positions rather than a single one simply because more chances are available. The result in Point 2 is not so obvious. Both types of contest rank a single player at the first level, but while one type indiscriminately ranks all other players at the last level, the other type selects also a single player for the second level. A player with a relatively low effort is more likely to be ranked first in the first type of contest, while a player with relatively high effort in the second type. Intuitively, the higher the number of levels in the outcome, the more a contest is "precise"

in identifying the player with the highest effort.²² In a contest with three levels, a player with low effort is more likely to be ranked third or second, and a player with high effort to be ranked second or first.

Let us see how the results in Proposition 5.2 apply to our examples with three players. Consider player 1's probability of being ranked first in the single-winner and the multi-winner contests. For the single-winner contest this is the probability of ranking (1; 2; 3), and for the multi-winner contest this is the joint probability of rankings (1, 2; 3) and (1, 3; 2). By Proposition 5.2 player 1's probability of being ranked first is always higher in the multi-winner contest than in the single-winner one. Consider the strict-ranking contest. Player 1's probability of being ranked first in this contest is the joint probability of (1; 2; 3) and (1; 3; 2). For $n = 3$, a strict ranking can be seen as a ranking with one player at first level, one player at second level and $n - 2$ players at third level. Then, by Proposition 5.2 player 1's probability of being ranked first is higher (lower) in the single-winner contest than in the strict ranking contest if and only if $f(x_1)$ is lower (higher) than the opponents' average $(f(x_2) + f(x_3))/2$.

5.3 Players' incentives in contests with various structures

In this Section we study contest games where players compete for some prizes by spending costly efforts. In Proposition 5.3 we study players' incentives in contests with various types of outcome. More specifically, we compare equilibrium efforts in single-winner, multi-winner and strict-ranking contests, establishing which types induce the highest efforts. While all these contests are organized in a single stage, in Proposition 5.4 we analyze contests with two stages and compare their equilibrium efforts with the ones of single-stage contests. Two-stage contests are studied under different assumptions on the timing of players' choices of efforts. More specifically, we analyze two-stage contests where all efforts are chosen in the first stage, and two-stage contests where efforts are chosen in both stages. Propositions 5.3 and 5.4 summarize results which are stated in more detail in their proofs in the Appendix.

Contests with various types of outcome

The outcome of a contest is a ranking of players. For any level $l \in N$ in the outcome, a player ranked at level l receives a prize $v_l \geq 0$. We assume $v_l \geq v_{l+1}$ for all $l \in N$ and $v_l > v_{l+1}$ for some $l \in N$. Players are risk neutral and efforts' costs are linear and sunk. Then, a player $i \in N$'s payoff is

$$\pi_i(x) = \sum_{l=1}^n \sum_{r \in R_{i,l}(t)} p(r, x) v_l - x_i$$

where $R_{i,l}(t)$ is the subset of $R(t)$ which includes any ranking r such that $r(i) = l$, for any $l \in N$. We define the *average prize* as $\bar{v} := (\sum_{l \in N} v_l)/n$ and the *average level* as $\bar{l} := (\sum_{l \in N} l)/n$, which is equal to $(n + 1)/2$ by Gauss formula.

Proposition 5.3 *Take a single-winner contest awarding a prize to the winner and nothing to others. In a symmetric interior equilibrium total efforts are*

²²This is not the case in strict-ranking contests with success function of Clark and Riis (1996). More precisely, with their success function a player's probability of being ranked first is always equal to the player's probability of being ranked first in a single-winner contest with same effort profile.

1. higher than the ones in the symmetric equilibrium of any multi-winner contest which equally shares the same prize among $m \geq 2$ winners and gives nothing to others.
 - (a) The equilibrium efforts of this multi-winner contest are equal to the ones of multiple separated single-winner contests, each with n/m players and each awarding one of these shares to its winner.
2. lower than the ones in the symmetric equilibrium of a strict-ranking contest awarding the same prize to the player ranked first and nothing to others.
 - (a) For any prize profile $(v_l)_{l \in N}$ the equilibrium efforts of this strict-ranking contest satisfy

$$\frac{f(x_i)}{f'(x_i)} = \bar{v} - \frac{1}{n} \sum_{l=1}^n l v_l \text{ for any } i \in N.$$

- (b) For any level $l \in N$, increasing the value of prize v_l induces higher (lower) equilibrium efforts if and only if l is smaller (larger) than the average level. Given a budget constraint $\sum_{l \in N} v_l \leq V$ equilibrium efforts are maximized by giving the whole budget to the player ranked first and nothing to others.

Proof: See Appendix.

Let us briefly discuss these results. Point 1 is intuitive and related to Point 1 of Proposition 5.2, which argues that it is always easier to be ranked first in a multi-winner contest than in a single-winner one. Point 1.a can be understood as a consequence of pair-swap consistency, which says that the relative performance of any pair of players should depend only on their relative efforts.²³ The result in Point 2 is not obvious, but it can be understood by referring to Point 2 in Proposition 5.2, which suggests that contests with more levels are more precise, hence more competitive.²⁴ In Point 2.a we derive a simple formula for equilibrium efforts of strict-ranking contests, with straightforward implications on maximization of equilibrium efforts in Point 2.b.²⁵ The intuition for Point 2.b is that competition is enhanced by rewarding best performing players and punishing worst performing ones.

Two-stage contests

A *two-stage contest* is a game where players compete for a single prize by spending costly efforts and the competition is organized in two stages. In the first stage all players engage in a multi-winner contest, which selects a set $M \subset N$ of winners.²⁶ The set of all possible sets of winners is denoted by \mathcal{M} . Only players in M access the second stage, where they engage in a single-winner contest where the winner receives a prize $v_1 > 0$ and the others nothing. We will often refer to the winner of the second stage as the winner of the two-stage contest.

²³Wärneryd (2001) already showed that a single-winner contest awarding a unique prize v_1 induces higher efforts than m symmetric separated single-winner contests each with n/m players and each awarding a prize v_1/m . The results in Points 1 and 1.a of Proposition 5.3 complete this analysis by comparing these contests with a multi-winner contest with same players and prizes.

²⁴This result is different from the analysis of strict-ranking contests with success function of Clark and Riis (1996). More specifically, a strict-ranking contest with success function of Clark and Riis (1996) which gives a unique first prize always induces the same equilibrium efforts of a single-winner contest awarding the same prize to the winner and nothing to others.

²⁵The study of equilibrium efforts in strict-ranking contests with other success functions has often been analytically demanding. See Clark and Riis (1996, 1998c) and Fu and Lu (2009, 2012).

²⁶The cardinality $m \geq 2$ of the set of winners fully determines the type of the contest in the first stage.

Notice that there is a different second-stage contest for each possible set of winners $M \in \mathcal{M}$ of the first stage. A two-stage contest has *contingent expenditures* when each player $i \in N$ spends a different effort in each stage. Denote by x_i^1 player $i \in N$'s effort in the first stage, and by x_i^M its effort in the second stage for a given $M \in \mathcal{M}$. For any $M \in \mathcal{M}$, if $i \notin M$ then $x_i^M = 0$. Call $x^1 \in X$ a first stage effort profile, and $x^M \in X$ a second stage effort profile for any $M \in \mathcal{M}$. For each set of second stage participants $M \in \mathcal{M}$, a player $i \in M$'s second stage payoff is

$$\pi_i^M(x^M) = p^M(i, x^M)v_1 - x_i^M$$

where $p^M(i, x^M)$ denotes the probability of the ranking where player i is ranked first. Given a second stage effort profile $x^M \in X$ for each $M \in \mathcal{M}$, a player $i \in N$'s first stage payoff is

$$\pi_i^1(x^1) = \sum_{M \in \mathcal{M}_i} p(M, x^1)\pi_i^M(x^M) - x_i^1$$

where \mathcal{M}_i is the set of all sets of winners $M \in \mathcal{M}$ where $i \in M$ and $p(M, x^1)$ denotes the probability of the ranking where players in M are ranked first.

A two-stage contest has *generic expenditures* when a player $i \in N$ makes a unique expenditure y_i in the first stage, which determines the first stage's effort and the second stage's effort for any $M \in \mathcal{M}$ as $x_i^1 = x_i^M = y_i$. A player $i \in N$'s payoff is

$$\pi_i(y) = \sum_{M \in \mathcal{M}_i} p(M, y)p^M(i, y)v_1 - y_i .$$

Proposition 5.4 *Take a single-winner contest awarding a prize to the winner and nothing to others. In a symmetric interior equilibrium its total efforts are*

1. *lower than the ones in the symmetric equilibrium of a two-stage contest with generic expenditures awarding the same prize to the winner and nothing to others. The equilibrium efforts of this two-stage contest are maximized by eliminating (the closest natural number to) $n - \sqrt{n}$ players in the first stage.*
2. *higher than the ones in the symmetric subgame perfect equilibrium of a two-stage contest with contingent expenditures awarding the same prize to the winner and nothing to others, given f is a power function.²⁷*

Proof: See Appendix.

Let us comment these results. Point 1 says that a contest which awards a prize in two stages induces higher efforts than the one which awards it in a single stage if its expenditures are generic, while Point 2 says that the opposite holds if the contest has contingent expenditures. Notice that a two-stage contest implicitly ranks players in three levels: first stage losers, second stage losers and the second stage winner. If efforts are generic, a two-stage contest is analogous to a single-stage contest with three levels. Then, the intuition for Point 1 may

²⁷A function f is a *power function* if it takes the form $f(x_i) = (x_i + \beta)^\alpha$ for some $\alpha, \beta > 0$. Notice that a power function is always positive, increasing and twice differentiable. A power function is weakly concave if and only if $\alpha \leq 1$. The power function is the most popular functional form in the contests' literature and the similar case with $\beta = 0$ is axiomatically characterized in Skaperdas (1996).

follow from Point 2 of Proposition 5.3, which suggests that adding levels in the outcome induces higher efforts. The intuition for Point 2 is clearer, as with contingent efforts a player spends in the second stage only if it has passed the first stage, hence some players do not exert any effort in the second stage.²⁸

6 Appendix

Proof of Theorem 4.1

It is easy to verify that (1) satisfies pair-swap consistency. Moreover, for $n = 2$ the only type of ranking in T is single-winner, and pair-swap consistency directly defines p for any $r \in R(t)$ and $x \in X$. Let us show that pair-swap consistency requires p to take the functional form in (1) for all $n \geq 3$, $t \in T$, $r \in R(t)$ and $x \in X$. Given any $n \geq 3$, $t \in T$ and $x \in X$, pair-swap consistency requires

$$\frac{p(r, x)}{p(r_{i,j}, x)} = \frac{f(x_i)}{f(x_j)} \quad (2)$$

for any $r \in R(t)$ and pair of players $i, j \in N$ ranked at consecutive levels ($r(j) = r(i) + 1$). Such players always exist since $n \geq 3$ and there are at least two levels in a ranking. Given any $n \geq 3$, consider types $t \in T$ with only *two levels*. Among these types, let us focus first on the ones with at least two players at the second level. For any of these types, without loss of generality consider a ranking $r \in R(t)$ such that $r = (1, \dots; 2, 3, \dots)$. Take any $x \in X$. By (2) $p(r, x)/p(r_{1,2}, x) = f(x_1)/f(x_2)$, hence there exists a function $F(x) \neq 0$ such that $p(r, x) = f(x_1)F(x)$ and $p(r_{1,2}, x) = f(x_2)F(x)$. By the same argument, as $p(r, x)/p(r_{1,3}, x) = f(x_1)/f(x_3)$, there exists $F'(x) \neq 0$ such that $p(r, x) = f(x_1)F'(x)$ and $p(r_{1,3}, x) = f(x_3)F'(x)$. As $p(r, x) = f(x_1)F(x) = f(x_1)F'(x)$, we necessarily have $F(x) = F'(x)$, hence

$$p(r, x) = f(x_1)F(x), p(r_{1,2}, x) = f(x_2)F(x) \text{ and } p(r_{1,3}, x) = f(x_3)F(x). \quad (3)$$

Among the types with only two levels, let us now focus on the ones with at least two players at the first level. For any of these types, without loss of generality consider a ranking $r \in R(t)$ such that $r = (1, 2, \dots; 3, \dots)$. Take any $x \in X$. By (2) $p(r, x)/p(r_{1,3}, x) = f(x_1)/f(x_3)$, hence there exists $F(x) \neq 0$ such that $p(r, x) = f(x_1)F(x)$ and $p(r_{1,3}, x) = f(x_3)F(x)$. Moreover, $p(r, x)/p(r_{2,3}, x) = f(x_2)/f(x_3)$, hence there exists $F'(x) \neq 0$ such that $p(r, x) = f(x_2)F'(x)$ and $p(r_{2,3}, x) = f(x_3)F'(x)$. We now have obtained $p(r, x) = f(x_1)F(x) = f(x_2)F'(x)$. Then, there exists $F''(x) \neq 0$ such that $p(r, x) = f(x_1)f(x_2)F''(x)$, and we necessarily have

$$p(r, x) = f(x_1)f(x_2)F''(x), p(r_{1,3}, x) = f(x_2)f(x_3)F''(x)$$

²⁸This result relates to previous studies of multiple-stage contests with contingent expenditures and different success functions. Clark and Riis (1998c) and Fu and Lu (2012) show that adding a stage to a contest with contingent expenditures always induces higher efforts. As argued in Point 2 of Proposition 5.4 the opposite holds with our success function. The reason is not obvious. Here is a possible interpretation. As they focus on success functions of strict-ranking type only, adding a stage to a contest implicitly adds levels in the outcome. Then, the increase in equilibrium efforts may be due to adding levels in the outcome.

$$\text{and } p(r_{2,3}, x) = f(x_1)f(x_3)F''(x). \quad (4)$$

Given any $x \in X$, by iteratively applying the arguments which brought to (3) and (4) one can show that for every type of ranking t with only two levels there exists a function $F_t(x) \neq 0$ such that $p(r, x) = \prod_{i \in A_r} f(x_i)F_t(x)$ for each $r \in R(t)$, where $A_r \subset N$ is the set of all payers ranked at first level in r . Notice that this expression can also be written as $p(r, x) = \prod_{i \in N} f(x_i)^{-r(i)}G_t(x)$, where $G_t(x) = F_t(x) \prod_{i \in N} f(x_i)^2$. By the definition of contest success function, the sum of the probabilities of all rankings in $R(t)$ is 1, hence

$$G_t(x) = \sum_{r \in R(t)} \prod_{i \in N} f(x_i)^{-r(i)}.$$

It follows that $p(r, x)$ must be as in (1) for any $n \geq 3$, $x \in X$, type t with only two levels and ranking $r \in R(t)$.

Given any $n \geq 3$ and type $t \in T$ with *at least three levels*, consider any ranking $r \in R(t)$. Take any triple of players $i, j, h \in N$ ranked $r(h) = r(j) + 1 = r(i) + 2$. Given r , consider the ranking $r' \in R(t)$ where i, j, h are ranked $r'(h) = r(i)$, $r'(j) = r(h)$ and $r'(i) = r(j)$ and all other players are ranked as in r . With some abuse of notation, these two rankings are represented by $r = (i \dots; j \dots; h \dots)$ and $r' = (h \dots; i \dots; j \dots)$. Take any $x \in X$. As the pair-swap ranking $r_{j,h} = (h \dots; j \dots; i \dots)$ is equal to the pair-swap ranking $r'_{i,h}$, we have

$$\frac{p(r, x)}{p(r', x)} = \frac{p(r, x)}{p(r_{j,h}, x)} \frac{p(r'_{i,h}, x)}{p(r', x)}.$$

By (2), $p(r, x)/p(r_{j,h}, x) = f(x_j)/f(x_h)$ and $p(r', x)/p(r'_{i,h}, x) = f(x_i)/f(x_h)$, hence

$$\frac{p(r, x)}{p(r', x)} = f(x_i)f(x_j)f(x_h)^{-2}. \quad (5)$$

Notice that $r'(i) - r(i) = 1$, $r'(j) - r(j) = 1$ and $r'(h) - r(h) = -2$. Then, the RHS of (5) can be written as $\prod_{k \in \{i,j,h\}} f(x_k)^{r'(k) - r(k)}$. As the pair-swap ranking $r_{i,h}$ is equal to the pair-swap ranking $r'_{i,j}$, we have

$$\frac{p(r, x)}{p(r_{i,h}, x)} = \frac{p(r, x)}{p(r', x)} \frac{p(r', x)}{p(r'_{i,j}, x)}.$$

By (2) and (5), we have

$$\frac{p(r, x)}{p(r_{i,h}, x)} = f(x_i)^2 f(x_h)^{-2} \quad (6)$$

Again, notice that $r_{i,h}(i) - r(i) = 2$, $r_{i,h}(j) - r(j) = 0$ and $r_{i,h}(h) - r(h) = -2$, and the RHS of (6) can be written as $\prod_{k \in \{i,j,h\}} f(x_k)^{r_{i,h}(k) - r(k)}$. By iteratively applying the arguments which led to (5) and (6), we obtain

$$\frac{p(r, x)}{p(r', x)} = \prod_{i \in N} f(x_i)^{r'(i) - r(i)} \text{ for each pair of rankings } r, r' \in R(t). \quad (7)$$

By (7), there must exist a function $G_t(x) \neq 0$ such that

$$p(r, x) = \prod_{i \in N} f(x_i)^{-r(i)} G_t(x) \text{ for each } r \in R(t).$$

By the definition of contest success function the sum of the probabilities of all rankings in $R(t)$ is 1, hence $G_t(x) = \sum_{r \in R(t)} \prod_{i \in N} f(x_i)^{-r(i)}$. It follows that $p(r, x)$ must be as in (1) for any $n \geq 3$, $x \in X$, $t \in T$ and $r \in R(t)$. □

Proof of Proposition 5.1

Given a function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_{++}$, for any $t \in T$ and $x \in X$, the probability of a ranking $r' \in R(t)$ is

$$p(r', x) = \frac{\prod_{i \in N} f(x_i)^{-r'(i)}}{\sum_{r \in R(t)} \prod_{i \in N} f(x_i)^{-r(i)}}.$$

Assume f twice differentiable, increasing ($f' > 0$) and weakly concave ($f'' \leq 0$). The first derivative of $p(r', x)$ with respect to x_i is

$$\frac{\partial}{\partial x_i} [p(r', x)] = \frac{f'(x_i)}{f(x_i)} p(r', x) [\rho(i, x) - r'(i)].$$

For any $x \in X$, $\partial [p(r', x)] / \partial x_i \geq 0$ if and only if $r'(i) < \rho(i, x)$, hence $p(r', x)$ is increasing in x_i if r' ranks player i above its expected level in the outcome, while $p(r', x)$ is decreasing otherwise. Consider any type of ranking $t \in T$ with only two levels. As $r(i) \in \{1, 2\}$ for any $r \in R(t)$, then $\rho(i, x) \in [1, 2]$ for any $x \in X$. Then, as $\rho(i, x) \geq 1$ for any $x \in X$, the probability $p(r', x)$ is always increasing in x_i for any $r' \in R(t)$ where $r'(i) = 1$. Similarly, $p(r', x)$ is decreasing in x_i for any $r' \in R(t)$ where $r'(i) = 2$.

Let us consider concavity. The second derivative of $p(r', x)$ with respect to x_i is

$$\begin{aligned} \frac{\partial^2}{\partial x_i^2} [p(r', x)] &= H(x_i) p(r', x) [r'(i) - \rho(i, x)] + \\ &+ \left[\frac{f'(x_i)}{f(x_i)} \right]^2 p(r', x) [d_i(r', x) - \delta_i(x)] \end{aligned} \quad (8)$$

where $H(x_i) = (f'(x_i)/f(x_i))^2 - f''(x_i)/f(x_i)$. The RHS of (8) is the sum of two parts. As $H(x_i) > 0$ for any $x_i \geq 0$, the first part is negative if and only if $r'(i) < \rho(i, x)$, hence $p(r', x)$ tends to be concave when increasing, and convex when decreasing. The second part is negative if and only if $d_i(r', x) < \delta_i(x)$. Then, $r'(i) < \rho(i, x)$ and $d_i(r', x) < \delta_i(x)$ are sufficient conditions for the concavity of $p(r', x)$ in x_i , while $r'(i) > \rho(i, x)$ and $d_i(r', x) > \delta_i(x)$ are sufficient conditions for its convexity.

Consider any type of ranking t with only two levels. For each player $i \in N$, take any $r \in R(t)$ where i is ranked first ($r(i) = 1$). By (8) $\rho(i, x)^2 \leq 3\rho(i, x) - 2$ is a sufficient condition for $p(r, x)$ being concave in x_i for any $x \in X$. As rankings have only two levels, we have $\rho(i, x) \in [1, 2]$. The LHS and RHS of the condition are equal to each other at $\rho(i, x) = 1$ and $\rho(i, x) = 2$. Moreover the LHS is convex in $\rho(i, x)$, while the RHS is linear. It follows that the LHS is smaller than the RHS for any $\rho(i, x)$ and $p(r, x)$ is concave in x_i for any $x \in X$. By similar arguments, one can show that for any type $t \in T$ with only two levels $p(r, x)$ is convex in x_i for any $x \in X$ and $r \in R(t)$ where $r(i) = 2$.

□

Proof of Proposition 5.2

Without loss of generality, let us focus on the probabilities of player 1 being ranked first. The probability of player 1 being ranked first in a single-winner contest is $f(x_1)/[\sum_{i \in N} f(x_i)]$ for some f positive and increasing. Consider a multi-winner contest with $m \geq 2$ winners. Call \mathcal{M} the set of all $M \subset N$ of cardinality m . \mathcal{M} is the set of all sets of winners. For any player $i \in N$, \mathcal{M}_i is the set of all sets of winners $M \in \mathcal{M}$ where $i \in M$. With some abuse of notation, for each $M \in \mathcal{M}$ denote by $p(M, x)$ the probability of the ranking $r \in R(t)$ where $r(i) = 1$ for any $i \in M$. Then, the probability of 1 being ranked first is $\sum_{M \in \mathcal{M}_1} p(M, x)$. For any $x \in X$, player 1 is more likely to be ranked first in the multi-winner contest than in the single-winner contest if and only if

$$\frac{\sum_{M \in \mathcal{M}_1} \prod_{i \in M} f(x_i)}{\sum_{M \in \mathcal{M}} \prod_{j \in M} f(x_j)} > \frac{f(x_1)}{\sum_{j \in N} f(x_j)}$$

After few manipulations this condition becomes

$$\sum_{M \in \mathcal{M}_1} \prod_{i \in M \setminus \{1\}} f(x_i) \left[\sum_{j \in N} f(x_j) \right] > \sum_{M \in \mathcal{M}_1} \prod_{i \in M \setminus \{1\}} f(x_i) \left[\sum_{j \notin M} f(x_j) + f(x_1) \right]$$

which is always true.

Consider a contest of a type with one player at the first level, one player at the second level and $n - 2$ players at the third level. For any $x \in X$, player 1's probability of being ranked first in this contest is

$$\frac{\sum_{j \neq 1} f(x_1)^2 f(x_j)}{\sum_{i \in N} \sum_{j \neq i} f(x_i)^2 f(x_j)}.$$

Then, comparing this probability with $f(x_1)/[\sum_{i \in N} f(x_i)]$, one can show that player 1 is more likely to be ranked first in the single winner contest if and only if

$$f(x_1) < \frac{\sum_{j \neq 1} f(x_j)^2 (\sum_{k \neq 1, j} f(x_k))}{\sum_{j \neq 1} f(x_j) (\sum_{k \neq 1, j} f(x_k))} \text{ for any } x \in X.$$

□

Proof of Points 1 and 1.a of Proposition 5.3

The players in the set $K = \{1, \dots, k\}$ ($k \geq 2$) engage in a single-winner contest which gives a prize v_1 to the winner and nothing to others. By Proposition 5.1 the probability of a ranking where player $i \in K$ is ranked first is increasing and concave in x_i for any $x \in X$, hence player i 's payoff is concave in x_i . One can show that for any f where $f(0)/f'(0) < (k-1)v_1/k^2$ there exists a unique equilibrium where each player $i \in K$ exerts effort x_i^* such that

$$\frac{f(x_i^*)}{f'(x_i^*)} = \frac{(k-1)v_1}{k^2}. \quad (9)$$

Consider $m \geq 2$ single-winner contests, each equivalent to the one above. There are m groups of k players, hence $n = m \cdot k$ players in total. The set of all players is $N = \{1, \dots, n\}$. As the k players of each group compete only within their group, equilibrium efforts are as in (9).

Suppose now that the players in N do not compete only within their group, but contend the m prizes in a grand contest all against all. Consider a grand contest of single-winner type, where the winner gets all the m prizes (a prize of value $m \cdot v_1$) and others nothing. One can show that for any f where $f(0)/f'(0) < (n-1)mv_1/n^2$ there exists a unique equilibrium where each player $i \in N$ exerts effort x_i^* such that

$$\frac{f(x_i^*)}{f'(x_i^*)} = \frac{(n-1)mv_1}{n^2}. \quad (10)$$

Suppose now that in the grand contest each player $i \in N$ can obtain *at most one* of the m prizes. The grand contest is a multi-winner contest with n players and m winners ($t(1) = m$ and $t(2) = n - m$), where each winner gets a prize v_1 and others nothing. Let \mathcal{M} be the set of all subsets of N of cardinality m , representing the set of all possible sets of winners $M \in \mathcal{M}$. For any $x \in X$ and $M \in \mathcal{M}$, the probability that M is the set of winners is

$$p(M, x) = \frac{\prod_{i \in M} f(x_i)}{\sum_{M' \in \mathcal{M}} \prod_{i \in M'} f(x_i)}$$

where with some abuse of notation $p(M, x)$ represents the probability of the ranking $r \in R(t)$ where $r(i) = 1$ for each $i \in M$. The payoff of a player $i \in N$ is

$$\pi_i(x) = \sum_{M \in \mathcal{M}_i} p(M, x)v_1 - x_i$$

where \mathcal{M}_i is the set of all $M \in \mathcal{M}$ where $i \in M$. Assume f increasing, concave and twice differentiable. By Proposition 5.1 the probability of a ranking where a player $i \in N$ is ranked first is increasing and concave in x_i . Then, player i 's payoff is concave in x_i and an interior best reply is identified by the first order condition

$$\sum_{M \in \mathcal{M}_i} \frac{\partial}{\partial x_i} p(M, x)v_1 = 1.$$

It can be shown that for any f where $f(0)/f'(0) < m(n-m)v_1/n^2$ there exists a symmetric interior equilibrium where the effort of each player $i \in N$ is such that

$$\frac{f(x_i^*)}{f'(x_i^*)} = \frac{m(n-m)v_1}{n^2} \quad (11)$$

It is easy to verify that the RHS of (10) is always larger than the RHS of (11), hence the single-winner grand contest always induces higher efforts than the multi-winner grand contest. Moreover, as $n = m \cdot k$, the RHS of (11) is equal to the RHS of (9), hence the multi-winner grand contest induces the same efforts as the separated single-winner contests. Corollary 6.1 summarizes the results.

Corollary 6.1 *Suppose f is positive, increasing, weakly concave and twice differentiable. For some f , there exists an equilibrium of the multi-winner grand contest where efforts are equal to the ones in the unique equilibrium of single-winner contests with same prizes. Moreover these efforts are smaller than the ones in the unique equilibrium of the single-winner grand contest which gives all prizes to the winner.*

□

Proof of Points 2, 2.a and 2.b of Proposition 5.3

Consider a strict-ranking contest with any prize profile $(v_l)_{l \in N}$. If there exists an interior equilibrium, the best reply of each player $i \in N$ is identified by the first order condition

$$\sum_{l=1}^n \sum_{r \in R_{i,l}(t)} \frac{\partial}{\partial x_i} [p(r, x)] v_l = 1 .$$

If the equilibrium is symmetric, the equilibrium effort x_i^* of each player $i \in N$ is such that

$$\frac{f(x_i^*)}{f'(x_i^*)} = \frac{1}{n} \sum_{l=1}^n [\bar{l} - l] v_l \quad (12)$$

and can also be written as $f(x_i^*)/f'(x_i^*) = \bar{l}\bar{v} - 1/n \sum_{l=1}^n l v_l$. The RHS of (12) is a weighted average of the prizes. Notice that the weight of prize v_l is positive if and only if l is smaller than the average level \bar{l} . Then, given a prize profile $(v_l)_{l \in N}$, increasing a prize v_l induces higher efforts if and only if $l < \bar{l}$.

Notice that in the RHS of (12) the weight of a prize v_l is always larger than the weight of v_{l+1} for any $l < n$. Then, given a limited budget $\sum_{l \in N} v_l \leq V$, equilibrium efforts are maximized by giving a unique large first prize $v_1 = V$. By (12), if $v_1 = V$, each player exerts effort such that $f(x_i^*)/f'(x_i^*) = (n-1)V/(2n)$. Consider the single-winner contest which gives a unique prize V to the winner. It is easy to verify that in a symmetric interior equilibrium this contest induces efforts such that $f(x_i^*)/f'(x_i^*) = (n-1)V/n^2$. It follows that, in a symmetric interior equilibrium, the strict ranking contest with $v_1 = V$ always induces higher efforts than the single-winner contest with prize V . Corollary 6.2 summarizes the results.

Corollary 6.2 *Suppose f is positive, increasing, weakly concave and twice differentiable. In a symmetric interior equilibrium of a strict-ranking contest efforts are as in (12) and*

1. *given a prize profile $(v_l)_{l \in N}$, increasing v_l induces higher efforts if and only if $l < \bar{l}$ for any $l \in N$;*
2. *given a budget $\sum_{l \in N} v_l \leq V$, equilibrium efforts are maximized by the prize profile $v_1 = V$ and $v_l = 0$ for all $l \geq 2$. These efforts are higher than the ones in the symmetric interior equilibrium of a single-winner contest which gives V to the winner and nothing to others.*

Let us see an example where the effort profile in Corollary 6.2 is an equilibrium. Let $N = \{1, 2, 3\}$. The set of possible outcomes is

$$R(t) = \{(1; 2; 3), (1; 3; 2), (2; 1; 3), (2; 3; 1), (3; 1; 2), (3; 2; 1)\}.$$

Without loss of generality, suppose $v_3 = 0$. Player 1's payoff is

$$\pi_1(x) = [p((1; 2; 3), x) + p((1; 3; 2), x)] v_1 + [p((2; 1; 3), x) + p((3; 1; 2), x)] v_2 - x_1$$

and similarly defined for other players. By (12) we know that, if a symmetric interior equilibrium exists for some f , equilibrium efforts are such that $f(x_i^*)/f'(x_i^*) = v_1/3$ for all $i \in N$, rankings' probabilities are $p(r, x^*) = 1/6$ for all $r \in R(t)$ and each player $i \in N$'s expected level in the outcome is $\rho(i, x^*) = 2$. Take any f where $f(0)/f'(0) < v_1/3$. By Proposition 5.1, for any $x \in X$, $r \in R(t)$ and $i \in N$, the probability $p(r, x)$ is increasing in x_i if and only if $r(i)$ is smaller than $\rho(i, x)$. Then, for any $r \in R(t)$ and $i \in N$, the probability $p(r, x)$ valued at x^* is increasing in x_i if $r(i) > 2$ and constant if $r(i) = 2$. Moreover, for any $r \in R(t)$ and $i \in N$ one can show that the probability $p(r, x)$ valued at x^* is concave in x_i if $r(i) \leq 2$. Then, for any $r \in R(t)$ and $i \in N$, the probability $p(r, x)$ valued at x^* is increasing and concave in x_i if $r(i) = 1$ and constant and concave in x_i if $r(i) = 2$. It follows that player i 's payoff at x^* is concave in x_i , and x^* is an equilibrium for some f . Corollary 6.3 summarizes the results.

Corollary 6.3 *For $n = 3$ the effort profile in Corollary 6.2 is an equilibrium for some f .*

□

Proof of Point 1 of Proposition 5.4

Consider a two-stage contest with generic expenditures which gives a prize v_1 to the final winner and nothing to others. If an interior equilibrium exists, the best reply of each player is identified by a first order condition. In a symmetric equilibrium, the efforts are such that

$$\frac{f(y_i^*)}{f'(y_i^*)} = \frac{v_1}{n} \left(2 - \frac{m}{n} - \frac{1}{m} \right). \quad (13)$$

By (15), a one-stage contest with prize v_1 induces efforts such that

$$\frac{f(y_i^*)}{f'(y_i^*)} = \frac{v_1}{n} \left(1 - \frac{1}{n} \right). \quad (14)$$

For any n and m the RHS of (14) is smaller than the RHS of (13). Then, any two-stage contest induces higher efforts than the one-stage contest. Let us find m which maximizes the efforts of a two-stage contest. It is easy to verify that the RHS of (13) is maximized at (the closest natural number to) $m^* = \sqrt{n}$, taking value (approximately)

$$\frac{f(y_i^*)}{f'(y_i^*)} = \frac{2v_1}{n} \left(1 - \frac{1}{\sqrt{n}}\right).$$

Corollary 6.4 summarizes the results.

Corollary 6.4 *Suppose f is positive, increasing, weakly concave and twice differentiable. In a symmetric interior equilibrium of a two-stage contest*

1. *efforts are always higher than the ones in the symmetric interior equilibrium of a one-stage contest with same prize profile.*
2. *equilibrium efforts are maximized when the contest eliminates (the closest natural number to) $n - \sqrt{n}$ players in stage-1;*

Let us show an example where the effort profile described in Corollary 6.4 is an equilibrium. Consider $n = 4$ and $m = \sqrt{n} = 2$. Without loss of generality, let us focus on player 1. The set of all sets of stage-1 winners to which 1 belongs is $\mathcal{M}_i = \{\{1, 2\}, \{1, 3\}, \{1, 4\}\}$. The payoff of player 1 is $\pi(y) =$

$$\begin{aligned} &= p(\{1, 2\}, y) p^{\{1,2\}}(1, y) v_1 + p(\{1, 3\}, y) p^{\{1,3\}}(1, y) v_1 + \\ &\quad + p(\{1, 4\}, y) p^{\{1,4\}}(1, y) v_1 - y_1. \end{aligned}$$

The payoff of 1 is concave in y_1 if $p(M, y) p^M(1, y)$ is concave for any $M \in \mathcal{M}_i$. It is easy to verify that the second derivative of $p(M, y) p^M(1, y)$ with respect to y_1 , when valuated at a symmetric effort profile, is always negative. Then, player $i \in N$'s payoff $\pi_i(y)$ is concave in y_i when valuated at y^* . Moreover, one can show that for any f where $f(0)/f'(0) < v_1/4$ the effort profile y^* is an equilibrium.

Corollary 6.5 *For $n = 4$ and $m = 2$ the effort profile in Corollary 6.4 is an equilibrium for some f .*

□

Proof of Point 2 of Proposition 5.4

Consider a two-stage contest with contingent expenditures which gives a prize v_1 to the final winner and nothing to others. By results in Proposition 5.1, for any $M \in \mathcal{M}$ a player $i \in M$'s stage-2 payoff is concave in x_i^M . Then, one can show that for any f where $f(0)/f'(0) < (m-1)v_1/m^2$ there exists an equilibrium of stage-2 contest where each player $i \in M$ exerts effort x_i^{M*} such that

$$\frac{f(x_i^{M*})}{f'(x_i^{M*})} = \frac{(m-1)v_1}{m^2} \tag{15}$$

and its stage-2 payoff is $\pi_i^M(x^{M*}) = v_1/m^2$. By backward induction, a player $i \in N$'s stage-1 payoff is

$$\pi_i^1(x^1) = \sum_{M \in \mathcal{M}_i} \frac{\prod_{i \in M} f(x_i^1)}{\sum_{M' \in \mathcal{M}} \prod_{i \in M'} f(x_i^1)} \pi_i^M(x^{M*}) - x_i^1$$

where \mathcal{M}_i is the set of all $M \in \mathcal{M}$ where $i \in M$. It can be shown that, by results in Proposition 5.1, a player $i \in N$'s stage-1 payoff is concave in x_i^1 . For any f where $f(0)/f'(0) < (n-m)v_1/(n^2m)$ there exists an equilibrium where each player $i \in N$ exerts effort x_i^{1*} such that

$$\frac{f(x_i^{1*})}{f'(x_i^{1*})} = \frac{(n-m)v_1}{n^2m} . \quad (16)$$

The sum of all efforts spent by all players in the two stages is $nx_i^{1*} + mx_i^{M*}$. As this sum cannot be computed from (15) and (16) for every function f , consider f such that $f(x_i^1) = (x_i^1 + \beta)^\alpha$, where $\alpha \in (0, 1]$ and $\beta > 0$. With this functional form the equilibrium efforts are

$$x_i^{1*} = \frac{\alpha(n-m)v_1}{n^2m} - \beta , x_i^{M*} = \frac{\alpha(m-1)v_1}{m^2} - \beta \quad (17)$$

for any $i \in N$ and $M \in \mathcal{M}$ and the sum of all efforts is

$$\frac{\alpha(n-1)v_1}{n} - (n+m)\beta . \quad (18)$$

Consider a one-stage contest where the n players contend all against all the same prize $v_1 > 0$. It is easy to verify that the total expenditures in the one-stage contest are $\alpha(n-1)v_1/n - n\beta$, hence always larger than the ones in (18) by the amount $m \cdot \beta$. Corollary 6.6 summarizes the results.

Corollary 6.6 *Suppose $f(x_i^1) = (x_i^1 + \beta)^\alpha$, where $\alpha \in (0, 1]$ and $\beta > 0$. For some α and β , there exists an equilibrium of a two-stage contest where*

1. *total efforts are always lower than the ones of a one-stage contest with same prize profile;*
2. *the limit of total efforts for $\beta \rightarrow 0$ is equal to total efforts of the one-stage contest.*

□

References

- Amann, E. and W. Leininger**, "Asymmetric all-pay auctions with incomplete information: the two-player case," *Games and Economic Behavior*, 1996, 14 (1), 1–18.
- Arbatskaya, M. and H. M. Mialon**, "Multi-activity contests," *Economic Theory*, 2010, 43 (1), 23–43.

- Baye, M.R., D. Kovenock, and C.G. De Vries**, “Rigging the lobbying process: an application of the all-pay auction,” *American Economic Review*, 1993, *83* (1), 289–294.
- , – , and – , “The solution to the Tullock rent-seeking game when $R > 2$: Mixed-strategy equilibria and mean dissipation rates,” *Public Choice*, 1994, *81* (3), 363–380.
- , – , and – , “The all-pay auction with complete information,” *Economic Theory*, 1996, *8* (2), 291–305.
- Beggs, S., S. Cardell, and J. Hausman**, “Assessing the potential demand for electric cars,” *Journal of Econometrics*, 1981, *17* (1), 1–19.
- Blavatsky, P.R.**, “Contest success function with the possibility of a draw: Axiomatization,” *Journal of Mathematical Economics*, 2010, *46* (2), 267–276.
- Bozbay, I. and A. Vesperoni**, “Alliance networks in conflict,” *Mimeo*, 2013.
- Che, Y.K. and I. Gale**, “Difference-form contests and the robustness of all-pay auctions,” *Games and Economic Behavior*, 2000, *30* (1), 22–43.
- Clark, D.J. and C. Riis**, “A multi-winner nested rent-seeking contest,” *Public Choice*, 1996, *87* (1), 177–184.
- and – , “Competition over more than one prize,” *American Economic Review*, 1998, *88* (1), 276–289.
- and – , “Contest success functions: an extension,” *Economic Theory*, 1998, *11* (1), 201–204.
- and – , “Influence and the discretionary allocation of several prizes,” *European Journal of Political Economy*, 1998, *14* (4), 605–625.
- Cornes, R. and R. Hartley**, “Asymmetric contests with general technologies,” *Economic Theory*, 2005, *26* (4), 923–946.
- Esteban, J. and D. Ray**, “Conflict and distribution,” *Journal of Economic Theory*, 1999, *87* (2), 379–415.
- Fu, Q. and J. Lu**, “The beauty of bigness: On optimal design of multi-winner contests,” *Games and Economic Behavior*, 2009, *66* (1), 146–161.
- and – , “Micro foundations of multi-prize lottery contests: a perspective of noisy performance ranking,” *Social Choice and Welfare*, 2011, pp. 1–21.
- and – , “The optimal multi-stage contest,” *Economic Theory*, 2012, pp. 1–32.
- , – , and **Z. Wang**, “Reverse nested lottery contests,” *Mimeo*, 2013.
- Konrad, K. A and D. Kovenock**, “Multi-battle contests,” *Games and Economic Behavior*, 2009, *66* (1), 256–274.
- Krishna, V. and J. Morgan**, “An analysis of the war of attrition and the all-pay auction,” *Journal of Economic Theory*, 1997, *72* (2), 343–362.
- Luce, R.D.**, *Individual Choice Behavior a Theoretical Analysis*, John Wiley and sons, 1959.
- McFadden, D.**, “Conditional logit analysis of qualitative choice behavior,” *Frontiers in Econometrics*, 1973, pp. 105–142.

- , “Modeling the choice of residential location,” *Transportation Research Record*, 1978, (673).
- **and P. Zarembka**, “Conditional logit analysis of qualitative choice behavior,” *Frontiers in Econometrics*, 1974, p. 105.
- Moldovanu, B. and A. Sela**, “The optimal allocation of prizes in contests,” *American Economic Review*, 2001, pp. 542–558.
- **and –** , “Contest architecture,” *Journal of Economic Theory*, 2006, 126 (1), 70–96.
- Münster, J.**, “Group contest success functions,” *Economic Theory*, 2009, 41 (2), 345–357.
- Okuguchi, K. and F. Szidarovszky**, “On the existence and uniqueness of pure Nash equilibrium in Rent-seeking games,” *Games and Economic Behavior*, 1997, 18, 135–140.
- Rai, B.K. and R. Sarin**, “Generalized contest success functions,” *Economic Theory*, 2009, 40 (1), 139–149.
- Siegel, R.**, “All-Pay Contests,” *Econometrica*, 2009, 77 (1), 71–92.
- Skaperdas, S.**, “Contest success functions,” *Economic Theory*, 1996, 7 (2), 283–290.
- Thomson, W.**, “Consistency and its converse: an introduction,” *Review of Economic Design*, 2011, 15 (4), 257–291.
- Vesperoni, A.**, “Dominance solvable contests,” *Mimeo*, 2013.
- Wärneryd, K.**, “Replicating contests,” *Economics Letters*, 2001, 71 (3), 323–327.
- Williams, H.C.W.L.**, “On the formation of travel demand models and economic evaluation measures of user benefit,” *Environment and Planning A*, 1977, 9 (3), 285–344.